


# WILSON'S STORM PETREL

*Oceanites oceanicus oceanicus*



## Regional Occurrence (BCRs 14, 30)

- Distribution: pelagic in region
- Habitats used: pelagic; breeding in burrows on islands and in coastal areas

NAWCP ND=1

## Regional Abundance

- 1980s population size: 1,000,000-10,000,000nb
- Historical population size: unknown
- % change in regional population: unknown

NAWCP PS=2; NAWCP PT=3

## Regional Threats

- Identified threats: unknown
- Potential threats: unknown

NAWCP TN=2

## Conservation Status: Abundance (N breeders) and Conservation Rank

Global	North America	BCR 14	BCR 30	Mid-Atlantic	S. New England	Gulf of Maine	Maritimes
?	50,000-100,000nb	?nb	?nb	?nb	?nb	?nb	?nb
IUCN- Not Evaluated	NAWCP- Not currently at risk	Not at risk (PI=?)	Not at risk (PI=?)	NJ- SZN; DE-SZN MD-SZN; VA-SN	NY-S?N; CT-no info; RI- SZN; MA-SN; VT- no info	MA-SN; NH-SZN; ME- S5N; NB-S5N, NS-S5N	QC-SN; NB-S5N PE-SZN; NS-S5N

## Conservation Needs: Sustainability:

**Foraging Habitat:** primarily a surface-feeder on zooplankton, euphausiids, and amphipods; feeds over the continental shelf, follows ships and whales; disperses widely in northern hemisphere, esp. abundant in Gulf Stream, can occur close to shore, even feeding along the surf line; **Issues:** ; **Population Goals:** ; **Habitat Goals:**